Bowel Preparation with Halflytely

- DO NOT EAT OR DRINK ANYTHING 4 HOURS BEFORE YOUR SCHEDULED ARRIVAL TIME. \square
- You must be accompanied by a friend or relative to drive and/or assist you home. We WILL NOT discharge you to a cab, bus or other transportation service without a responsible party with you. You may not drive for 12 hours following your procedure.

Materials You Will Need to Prepare For Your Procedure

- **Prescription** for the **HalfLytely** Bowel Prep which has been provided by our office \square
- Obtain the HalfLytely Bowel Prep from the Pharmacy
- **Clear Liquid Diet Instructions (see attached Clear Liquid Diet Sheet)**

Special Instructions Regarding Medications

- Do Not take Iron supplements for one week prior to the procedure date. Multivitamins are OK. Π
- **Stop** taking **Coumadin/Plavix days prior** to your scheduled procedure date.
- **Diabetic patients**:
- Do Not take Lasix, HCTZ or Dyazide (water pill) the day of the procedure
- Heart/Blood Pressure/ Thyroid/Seizure/Nerve Medications can be taken on the day of the procedure with a sip of water. \square
- Stop Effexor one day before the procedure
- Asthmatic patients should bring their INHALERS with them on the day of the procedure
- \square Other

The Day Before Your Exam

- DO NOT EAT ANY SOLID FOODS. YOU MAY ONLY HAVE CLEAR LIQUIDS UNTIL 4 HOURS BEFORE YOUR SCHEDULED ARRIVAL TIME.
- Take all of your normal prescription medications, unless advised otherwise.
- You may go to work, but be home by 6 pm to begin your bowel prep.
- **IMPORTANT NOTE:** follow the instructions below, NOT the instructions on the Prep's packaging. \square
- Prepare your **HalfLytely** solution in the morning with lukewarm water to the designated line and add a flavor packet. Shake well and refrigerate.

PM Procedure

- At 6 pm Take two (2) Bisacodyl Laxative tablets with water. \square
- At 6 am on the morning of your procedure. Begin drinking the HalfLytely laxative solution. Drink 8 ounces every 10-15 minutes – until the entire solution is consumed.

You must finish drinking this solution by 8 am.

It is best to drink each glass rapidly rather than sip it. Drinking the HalfLytely solution through a straw may also improve your tolerance for this bowel prep.

WITH AN AFTERNOON PROCEDURE, you may have CLEAR LIOUIDS up to 4 hours prior to your SCHEDULED **ARRIVAL TIME.** Other

- - ⇒ Questions about symptoms or medications call 484-356-1050 between 7:30 am- 3:30 pm
 - \Rightarrow General or reschedule questions call 610-431-3122
 - \Rightarrow If you reschedule, ask if you will need new prep instructions.

Patient: Print Name

Signature

Date